

FFH TESTING RESULTS




PATIENTS

25,725




TESTED

1,617



POSITIVE

247



NEGATIVE

1,370

POSITIVITY RATES



FFH Patients

7%


1,617 tests
247 positive



York Co.

7.8%

61,458 tests
4,606 positive




Pennsylvania

3.8%

1,942,956 tests
152,544 positive

FFH PATIENT POSITIVITY RATES - OVER TIME



SBIRT (Screening, Brief Intervention, & Referral to Treatment)

SBIRT is an approach to the delivery of early intervention and treatment to people with substance use disorders and those at risk of developing these disorders.

2019 FFH SBIRT data:



40,797

SBIRT Screenings

1,244

Warm-Handoffs

5,695

BH Visits

FATAL OPIOID OVERDOSES DURING COVID-19

(Confirmed or suspected)

8
January

17
February

24
March

14
April
(First Half)

Opioid ODs spiked in York County at beginning of pandemic

Find more info: <https://www.yorkdispatch.com/story/news/local/2020/04/16/recipe-disaster-opioid-ods-spike-york-county-amid-covid-19-pandemic/5137106002/>

NONFATAL OPIOID OVERDOSES DURING COVID-19

(Through April 16, 2020)

20
Nonfatal ODs

60
SUS Patients
Dropped out of care


From January 1, 2020 - April 16, 2020,
the number of Naloxone reversals were 140 more than all of 2019.

During late June, 40% of U.S. adults reported struggling with mental health or substance use[†]


ANXIETY/DEPRESSION SYMPTOMS



TRAUMA/STRESSOR-RELATED DISORDER SYMPTOMS



STARTED OR INCREASED SUBSTANCE USE



SERIOUSLY CONSIDERED SUICIDE[†]



[†]Based on a survey of U.S. adults aged ≥18 years during June 24-30, 2020

[†]In the 30 days prior to survey

For stress and coping strategies: bit.ly/dailylifecoping